

**Questions or comments?**  
 Call toll-free 1-888-670-6100  
 Monday- Friday 8am – 5pm EST  
 Retain original packaging for future reference.



**Distributed by:**  
 Premier® Dental Products Company, Plymouth Meeting, PA 19462  
 www.premusa.com Part # 9007285 [www.enamelon.com](http://www.enamelon.com)  
 United States Patent Numbers: US 5,993,784, US 5,711,936,  
 US 5,651,959 and other patents pending. Made in U.S.A.  
 Ultramulsion® is a registered trademark  
 of WhiteHill Oral Technologies, Inc.  
 NDC 48783-500-40 051438 Rev2 5P

**premier Enamelon®**

**Clinically Proven Active Ingredient for  
 Anticavity, Antigingivitis and Sensitive Teeth**

- Ultramulsion® is a patented saliva-soluble coating that moisturizes and soothes soft tissues of the mouth
- Clean Mint flavor enhanced with spilanthes, a natural herb which can boost saliva flow and flavor
- No Sodium Lauryl Sulfate (SLS), No Gluten, No Dyes

STRENGTHEN YOUR SMILE!

**ACP™**  
 Enamel Therapy

**premier**

**Enamelon®**

**Preventive Treatment Gel**  
 Clean Mint NET WT 4.0 OZ (113 g)

**ULTRAMULSION®**

Sample Inside! Enamelon® Fluoride Toothpaste NET WT 0.75 OZ (21.2 g)

**Enamelon® Preventive Treatment Gel**

Drug Facts	Purpose
<b>Active Ingredient</b> Stannous Fluoride 0.40% ..... Anticavity/Sensitivity/ (0.15% w/v fluoride ion)	<b>Purpose</b> Anticavity/Sensitivity/ Antigingivitis
<b>Uses</b> • Aids in the prevention of caries • Builds increasing protection against painful sensitivity of the teeth to cold, heat, acids, sweets or contact • Helps prevent gingivitis • Helps interfere with the harmful effects of plaque associated with gingivitis	

Drug Facts (continued)
<b>Warnings</b> <b>Keep out of the reach of children.</b> If more than used for brushing is accidentally swallowed, get medical help or contact a Poison Control Center right away. <b>Do not use</b> if gingivitis, bleeding, or redness persists for more than 2 weeks, see your dentist. See your dentist immediately if you have painful or swollen gums, pus from the gum line, loose teeth, or increasing spacing between the teeth. These may be signs or symptoms of periodontitis, a serious form of gum disease.

Drug Facts (continued)
Sensitive teeth may indicate a serious problem that may need prompt care by a dentist. <b>Stop use and ask a dentist</b> if the problem persists or worsens. Do not use this product for sensitivity relief longer than 4 weeks unless recommended by a dentist or doctor.
<b>Directions</b> Adults and children 6 years of age or older: Use once a day after brushing your teeth with a toothpaste. Apply the gel to your

Drug Facts (continued)
teeth and brush thoroughly. Allow the gel to remain on your teeth for 1 minute and then spit out. Do not swallow the gel. Do not rinse, eat or drink for 30 minutes after brushing. Instruct children under 12 years of age in the use of this product (to minimize swallowing). Supervise children as necessary until capable of using without supervision. Children under 6 years of age: Consult a dentist or doctor.
<b>Other information</b> This is an anticavity/fluoride preventive treatment gel, not a toothpaste. Read directions carefully before using. Products containing stannous fluoride may produce surface staining of the teeth. Adequate toothbrushing may prevent these stains which are not harmful or permanent and may be removed by your dentist. • Do not use if foil seal on tube is broken or missing • Keep tightly closed when not in use • Store at controlled room temperature
<b>Inactive ingredients</b> Acesulfame K, calcium sulfate, cocamidopropyl betaine, dimethicone, flavors, glycerin, lauroyl-sarcosine, monosodium phosphate, poloxamer 407, polyethylene glycol, pvm/ma copolymer, silica, sucralose

Enamelon® Fluoride Toothpaste	
Drug Facts	Purpose
<b>Active Ingredient</b> Stannous Fluoride 0.45% ..... Anticavity/Sensitivity/ (0.16% w/v fluoride ion)	<b>Purpose</b> Anticavity/Sensitivity/ Antigingivitis
<b>Uses</b> • Aids in the prevention of caries • Builds increasing protection against painful sensitivity of the teeth to cold, heat, acids, sweets or contact • Helps prevent gingivitis • Helps interfere with the harmful effects of plaque associated with gingivitis	
<b>Warnings</b> <b>Keep out of the reach of children under 6 years of age.</b> If more than used for brushing is accidentally swallowed, get medical help or contact a Poison Control Center right away. <b>Do not use</b> if gingivitis, bleeding, or redness persists for more than 2 weeks, see your dentist. See your dentist immediately if you have painful or swollen gums, pus from the gum line, loose teeth, or increasing spacing between the teeth. These may be signs or symptoms of periodontitis, a serious form of gum disease. Sensitive teeth may indicate a serious problem that may need prompt care by a dentist. <b>Stop use and ask a dentist</b> if the problem persists or worsens. Do not use this product for sensitivity relief longer than 4 weeks unless recommended by a dentist or doctor.	

Drug Facts (continued)
<b>Directions</b> Adults and children 2 years of age or older: Brush teeth thoroughly, preferably after each meal or at least twice a day, or as directed by a dentist or doctor. Instruct children under 6 years of age in good brushing and rinsing habits (to minimize swallowing). Supervise children as necessary until capable of using without supervision. Children under 2 years of age: Consult a dentist or doctor.
<b>Other information</b> • Products containing stannous fluoride may produce surface staining of the teeth. Adequate toothbrushing may prevent these stains which are not harmful or permanent and may be removed by your dentist. • Do not use if foil seal on tube is broken or missing • Keep tightly closed when not in use • Store at controlled room temperature
<b>Inactive ingredients</b> Glycerin, polyethylene glycol, silica, pvm/ma copolymer, cocamidopropyl betaine, calcium sulfate, poloxamer 407, dimethicone, lauroyl-sarcosine, TiO <sub>2</sub> , flavor, acesulfame k, monosodium phosphate, sucralose